


AD A093997

NOO

SURFACE CURRENTS

SOUTHWEST INDIAN OCEAN.



17 ~~SEPTEMBER 1977~~

REPRINTED 1980

APPROVED FOR PUBLIC RELEASE, DISTRIBUTION UNLIMITED

NAVAL OCEANOGRAPHIC OFFICE
NSTL STATION, MISSISSIPPI 39522

**Best
Available
Copy**

ABSTRACT

THIS ATLAS, AND THE SERIES OF WHICH IT IS A PART, IS COMPUTER GENERATED AND AUTOMATICALLY PLOTTED. IT MAKES AVAILABLE TO THE USER THE MOST RECENT SURFACE CURRENT DATA COLLECTED AND WILL BE UPDATED WHENEVER SUFFICIENT AMOUNTS OF DATA ARE ADDED TO THE DATA FILE. THIS AND THE OTHER ATLASES ARE BASED ON A VAST QUANTITY OF DATA AS COMPARED TO THE PREVIOUS MANUALLY-COMPILED EDITIONS PRINTED IN THE MID-THIRTIES.

THE SURFACE CURRENT INFORMATION IS BASED MAINLY ON SHIP DRIFT, WHICH IS THE DIFFERENCE BETWEEN THE DEAD RECKONING POSITION AND THE POSITION DETERMINED BY ANY TYPE OF NAVIGATIONAL FIX. THIS DIFFERENCE DESCRIBES THE DIRECTION AND SPEED OF THE CURRENT.

ACKNOWLEDGMENTS


Messrs. Raymond J. Beauchesne* and William E. Boisvert made major contributions to this atlas.


*Mr. Beauchesne presently is employed by the Bureau of Naval Personnel.

FOREWORD

THIS ATLAS, ONE IN A SERIES OF 43 REGIONAL SURFACE CURRENT ATLASES, IS PRODUCED TO FULFILL A NEED OF NAVY PLANNING STAFFS AND THE SCIENTIFIC AND INDUSTRIAL COMMUNITIES FOR THE LATEST AVAILABLE OCEAN SURFACE CURRENT DATA. THESE ATLASES ADD TO THE WEALTH OF NAUTICAL INFORMATION UPON WHICH OPERATIONAL PLANNING, NAVIGATIONAL SAFETY, AND SHIPPING ECONOMY DEPEND. RAPID PRODUCTION AND WIDE DISSEMINATION OF THIS ATLAS ARE MADE POSSIBLE BY THE LATEST COMPUTER TECHNIQUES.

THE CONSTANT IMPROVEMENT IN THE QUALITY OF SURFACE CURRENT DATA RECEIVED OVER THE YEARS IS MADE POSSIBLE LARGELY BY THE MORE THOROUGH REPORTS OF VOLUNTARY OBSERVERS IN RECENT YEARS. THE DEFENSE MAPPING AGENCY, THE OCEANOGRAPHIC OFFICE, AND THE USER OF THE ATLASES RELY ON THE PERSONAL OBSERVATIONS OF THE MAN WHO HAS "BEEN THERE." MARINERS, IN REPORTING THEIR OBSERVATIONS, RENDER A SERVICE NOT ONLY TO THEMSELVES BUT ALSO TO ALL "WHO GO DOWN TO THE SEA IN SHIPS." WITH THE ADVENT OF NUCLEAR POWER, ELECTRONIC NAVIGATION AIDS, AND 300,000-TON SHIPS, UP-TO-DATE, RAPIDLY DISSEMINATED ENVIRONMENTAL AND NAVIGATIONAL INFORMATION HAS BECOME INCREASINGLY IMPORTANT.


JOHN R. McDONNELL
Captain, U.S. Navy
Commander

Accession	
NTIS OF	
DTIC TAB	
Unannounced	
Justification	
By	
Distribution	
Availability	
Dist	
	

SURFACE CURRENT ATLASES


THIS SERIES OF COMPUTERIZED ATLASES REPLACES THE OLD HYDROGRAPHIC OFFICE ATLASES OF SURFACE CURRENTS (HOP 566, 568, 569, 570) WHICH WERE MANUALLY COMPILED FROM DATA OBTAINED DURING THE PERIOD 1903 - 1934. THESE NEW ATLASES CONFORM TO THE STANDARD NAVY OCEAN AREA AND REGION INDEX LIMITS SHOWN BELOW: e.g., NOO SP 1402-NP 10 COVERS NORTH PACIFIC REGION 10 EAST OF THE PHILIPPINES.

AS AMOUNTS OF NEW DATA

THESE GRAPHICS SHOW AREAS AS THE NORTH SEW CURRENTS ARE STRONGLY PREDICTABLE HOURLY CHANGES

RECENT IMPROVEMENTS IN THE DATA FILE ASSURE THE INCLUSION OF THE LATEST, HIGH QUALITY SURFACE CURRENT DATA AVAILABLE. THE FILE NOW CONTAINS MORE THAN 4,200,000 OBSERVATIONS AND A GENERAL UPDATE OF THE FILE WILL BE MADE

INDEX



SURFACE CURRENT ATLASES


THIS SERIES OF COMPUTERIZED ATLASES REPLACES THE OLD HYDROGRAPHIC OFFICE ATLASES OF SURFACE CURRENTS (HOP 566, 568, 569, 570) WHICH WERE MANUALLY COMPILED FROM DATA OBTAINED DURING THE PERIOD 1903 - 1934. THESE NEW ATLASES CONFORM TO THE STANDARD NAVY OCEAN AREA AND REGION INDEX LIMITS SHOWN BELOW: e.g., NOO SP 1402-NP 10 COVERS NORTH PACIFIC REGION 10 EAST OF THE PHILIPPINES.

AS AMOUNTS OF NEW DATA

THESE GRAPHICS Y
AREAS AS THE NORTH SEA
CURRENTS ARE STRONGLY
PREDICTABLE HOURLY CHA

RECENT IMPROVEMENTS IN THE DATA FILE ASSURE THE INCLUSION OF THE LATEST, HIGH QUALITY SURFACE CURRENT DATA AVAILABLE. THE FILE NOW CONTAINS MORE THAN 4,200,000 OBSERVATIONS AND A GENERAL UPDATE OF THE FILE WILL BE MADE

INDEX



Introduction

The Surface Current Data File, from which these atlases are derived, consists primarily of over four million ship set and drift observations. These data were collected by the Netherlands, Japan, Britain, France, and the United States. The file is supplemented by several thousand Geomagnetic Electrokinetograph (GEX) observations, mostly Japanese. The file spans the period from the early 1850's to the present. The earliest observations were collected by the Netherlands and Great Britain; those of the 1960's through the present are primarily United States data.


General Quality

The quality of this data file is considered high for this type of derived value. The data have been carefully screened for duplication; observations taken under adverse conditions (i.e. high winds and waves, time between observations greater than 12 hours) have been eliminated when warranted. Consideration was given to the reliability of the observer; doubtful shipboard computations of set and drift were edited; and observations with erroneous locations (mostly observations on land) have been eliminated. The accepted data are considered most useful when used collectively as in summaries where a number of observations show trends.

General Observation Technique

The set (direction) and drift (speed) are computed by the navigator from the difference between the dead reckoning (DR) position and the position determined by any type of navigational fix. The drift can be determined along any straight line track and includes all factors which cause changes in the DR position. When a fix is obtained, the current set (direction) is FROM the DR position TO the fix; the drift (speed) is equal to the distance in nautical miles between the DR and the fix, divided by the number of hours since the last fix. For successive observations, the TO POSITION of one observation becomes the FROM POSITION of the next observation.


Because the influence of current may vary along a ship's track, the MEAN POSITION of the track is assigned as the geographic location of the current observation. An example of a current computation is shown in the figure below.




EXAMPLE OF A SURFACE CURRENT (SHIP'S DRIFT) OBSERVATION

Data Presentation

The following legend shows two types of surface current presentations by 1° quadrangle, type 1 with 12 or more observations and type 2 with fewer than 12 observations. Where there are 11 or fewer observations within a 1° quadrangle, the total number of observations is shown within the 90° quadrant containing the observations.




If there are 12 or more observations by vector resultants as follows




(1) Persistent Current - 60 percent or more of all observations fall within a 45° sector of the 8-point compass.




(2) Prevailing - all observations fall within a 45° sector



(4) Bizonal Flow - Practically all observations are concentrated in opposite pairs of 45° sectors, and one pair contains at least 80 percent as many observations as the opposite pair. This generally indicates variability that occurs in zones of entrainment between opposing currents (see examples A and B, quadrangles 1, 2, and 3).



If there are 12 or more observations in a 1° quadrangle, the surface current is depicted by vector resultants as follows.





(1) Persistent Current - 60 percent or more of all observations fall within a 45° sector of the 8-point compass.

(2) Prevailing Current - 70 percent or more of all observations fall within two adjacent 45° sectors.


(3) Primary Current with Secondary Direction -
(a) Primary Current - 50 percent or more of all observations fall within three adjacent 45° sectors.

(b) Secondary Direction - 20 percent or more of all observations fall within a 45° sector, and the two resultant vector directions are separated by more than 90° of arc.



(4) Bizonal Flow - Practically all observations are concentrated in opposite pairs of 45° sectors, and one pair contains at least 80 percent as many observations as the opposite pair. This generally indicates variability that occurs in zones of entrainment between opposing currents (see examples A and B, quadrangles 1, 2, and 3).

(5) Variable Current - The 45° sector with most observations has less than 25 percent of all observations; direction is indeterminate.



consists primarily of collected by the is supplemented mostly Japanese. earliest observations 1960's through

ved value. The under adverse er than 12 hours) reliability edited; and ob- been eliminated. in summaries

on the difference y any type of track and includes ined, the current is equal to the umber of hours he observation


N POSITION of tion. An example

EM
D.
S
2
AN
O

1° quadrangle, ations. When ber of observa-

PIAL OBSERVATIONS).
ED FOR 1901 On
OBSERVATIONS.
FOR ALL OBSERVATIONS
Y DIRECTION, 232
DIRECTION, 0.6 KNOT
AST

1 2



SUMMER - JAN, FEB, MAR

45 E

50 E

55 E

60 E

65 E

70 E

29 S

30 S

35 S

40 S

45 S

49 S

45 E

50 E


55 E

60 E


65 E


70 E

SUMMER - JAN, FEB, MAR



AUTUMN - APR, MAY, JUN





WINTER - JUL. AUG. SEP


45 E

50 E

55 E

60 E

65 E



45 E

50 E


55 E

60 E


65 E


WINTER - JUL, AUG, SEP

70




SPRING - OCT, NOV, DEC





ANNUAL-JAN THROUGH DEC



DISTRIBUTION LIST

NAVY

CINCPACFLT (02M)
COMTHIRDFLT
COMSEVENTHFLT
COMSUBPAC
COMNAVAIRPAC
COMPATWINGSPAC
PATWINGSPAC DET ADAK
PATWING 1
COMNAVSURFPAC
DIRNAVOCEANMET
FLENUMEACEN
FLEWEACEN GUAM
FLEWEACEN PEARL
NAVWEASERVFAC SAN DIEGO
NAVWEASERVFAC YOKOSUKA
NWSO ASHEVILLE
NWSO ADAK
NWSO AGANA
NWSO ATSUGI
NWSO KADENA
NWSO MISAWA

OTHER GOVT.

NOAA/NODC
NOAA/NCC

PRIVATE & UNIVERSITIES

FLORIDA ST. UNIV.
LOUISIANA ST. UNIV.
MASS. INST. OF TECH
ORE. ST. UNIV.
TEXAS A&M UNIV.
UNIV. OF MIAMI
UNIV. OF R.I.
UNIV. OF WASH.
SCRIPPS INST OF OCEANOGRAPHY
WOODS HOLE OCEANOGRAPHIC INST.

FOREIGN

HYDROGRAPHER/R.A.N.
DEPT. TRANSPORTATION/AUSTRALIA

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER NOD SP1404-1N6	2. GOVT ACCESSION NO. AD-A093977	3. REPORT TYPE AND PERIOD COVERED Final	4. AUTHOR(s)
5. TITLE (and Subtitle) SURFACE CURRENTS SOUTHWEST INDIAN OCEAN	6. PERFORMING ORG. REPORT NUMBER	7. AUTHORING ORIGINATING ORGANIZATION	8. PERFORMING ORG. NAME AND ADDRESS Naval Oceanographic Office NSTL Station, MS 39522
9. CONTROLLING OFFICE NAME AND ADDRESS Naval Oceanographic Office	10. DISTRIBUTION STATEMENT (for this Report) Approved for public release; distribution unlimited.	11. REPORT DATE September 1977	12. NUMBER OF PAGES 14
13. DISTRIBUTION STATEMENT (for the abstract entered in Block 20, if different from Report)	14. SECURITY CLASS. (for this report)	15. SECURITY CLASS. (for this report)	16. SECURITY CLASS. (for this report)
17. SUPPLEMENTARY NOTES	18. ABSTRACT (Continue on reverse side if necessary and identify by block number) This atlas, and the series of which it is a part, is computer generated and automatically plotted. It makes available to the user the most recent surface current data collected and will be updated whenever sufficient amounts of data are added to the data file. This and the other atlases are based on a vast quantity of data as compared to the previous manually-compiled editions printed in the mid-thirties.	19. ABSTRACT (Continue on reverse side if necessary and identify by block number)	20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE
GPO 5102-010-66/1

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20. Cont.

The surface current information is based mainly on ship drift, which is the difference between the dead reckoning position and the position determined by any type of navigational fix. This difference describes the direction and speed of the current.



SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)